

A New Channel for the Detection of Planetary Systems Through Microlensing: I. Isolated Events Due to Planet Lenses

Rosanne Di Stefano¹, Richard A. Scalzo²

ABSTRACT

We propose and evaluate the feasibility of a new strategy to search for planets via microlensing observations. This new strategy is designed to detect planets in “wide” orbits, i.e., with orbital separation, a , greater than $\sim 1.5R_E$. Planets in wide orbits may provide the dominant channel for the discovery of planets via microlensing, particularly low-mass (e.g., Earth-mass) planets. This paper concentrates on events in which a single planet serves as a lens, leading to an isolated event of short-duration. We point out that a distribution of events due to lensing by stars with wide-orbit planets is necessarily accompanied by a distribution of shorter-duration events. The fraction of events in the latter distribution is proportional to the average value of $\sqrt{q}$, where q is the ratio between planet and stellar masses. The position of the peak or peaks also provides a measure of the mass ratios typical of planetary systems. We study detection strategies that can optimize our ability to discover isolated short-duration events due to lensing by planets, and find that monitoring employing sensitive photometry is particularly useful. If planetary systems similar to our own are common, even modest changes in detection strategy should lead to the discovery a few isolated events of short-duration every year. We therefore also address the issue of the contamination due to stellar populations of any microlensing signal due to low-mass MACHOs. We describe how, even for isolated events of short-duration, it will be possible to test the hypothesis that the lens was a planet instead of a low-mass MACHO, if the central star of the planetary system contributes a measurable fraction of the baseline flux.

Subject headings: – Gravitational lensing: microlensing, dark matter – Stars: planetary systems, luminosity function, mass function – Planets & satellites: general – Galaxy: halo – Methods: observational – Galaxies: Local Group.

¹ Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138; e-mail: rdistefano@cfa.harvard.edu

²Department of Physics, University of Chicago, Chicago, IL 60637; e-mail: rscalzo@rainbow.uchicago.edu

1. Introduction

When planets are discovered through monitoring programs similar to those presently being carried out, the distance to the lens will typically be on the order of kiloparsecs. Although some follow-up may be possible (DiStefano & Scalzo 1997, 1998, DiStefano 1998a,b), imaging, such as that planned to probe possible companions to nearby stars (Stahl & Sandler 1995; Bender & Stebbins 1996; Labeyrie 1996; Angel & Woolf 1997) will not be possible in the near future. Nevertheless, planets discovered as microlenses can play an important role in developing our understanding of planetary systems in our Galaxy and beyond. The reason for this is precisely because microlensing observations probe vast volumes of the Milky Way and other galaxies, such as the Magellanic Clouds and M31. Thus, microlensing provides a unique window for studying the statistics of planetary systems (numbers and properties) and their dependence on the local stellar environment. Microlensing searches complement velocity-based searches in another way as well. Particularly for the wide systems studied here for the first time, the microlensing searches can be effective for low-mass planets orbiting at low speeds; such planets cannot readily be detected via radial velocity methods.

The framework for the work to date on discovering planets through microlensing was established by Mao & Paczyński (1991) and Gould & Loeb (1992). These authors found that when the separation, a , between the star and planet is close to the Einstein radius, R_E , of the star ($0.8R_E \lesssim a \lesssim 1.5R_E$), a significant fraction of events ($\sim 5 - 20\%$) in which the star serves as a lens would be perturbed in a detectable way. This has been referred to as “resonant” lensing, both because the separation must be close to R_E in order for the signal to be detectable, and because the signal itself is sharp and distinctive. The detection strategy is to monitor light from an ongoing event at frequent intervals in order to observe a short-lived perturbation. With the analysis of more than 200 microlensing events reported to date, it is not clear whether any planetary-lens events have been discovered.³

³ One candidate for a resonant planetary lensing event has been suggested (Bennett *et al.* 1996a). There are, however, two reasons to be cautious about the interpretation of this event. First, the mass ratio derived from the binary-lens fit is ~ 0.043 , which is large enough to be consistent with lensing by a binary stellar system. Second, the degeneracy of the physical solution has not yet been worked out. The degeneracy may be of two types: (a) Other binary solutions with values of q differing from the one for this fit by as much as an order of magnitude may prove to be equally good fits (Di Stefano & Perna 1996); this needs to be systematically checked. (b) Other physical effects may prove to be important, making the determination of the system’s true parameters even more difficult. These effects include finite source size effects (Witt & Mao 1994; Witt 1995), finite lens-size effects (Bromley 1996), and blending (Di Stefano & Esin 1995, Kamionkowski 1995). (In fact blending has been a feature of every other binary-lens event; Udalski *et al.* 1994, Alard, Mao, and Guibert 1995, Alcock *et al.* 1997a, Bennett *et al.* 1996b.)

The purpose of this paper and its companion (DiStefano & Scalzo 1998) is to develop a complementary framework for the discovery of planets by microlensing. The basic idea is that planets located more than $\sim 1.5R_E$ from the central star can give rise to characteristic microlensing signatures that reveal evidence of their presence. The advantages of searching for planets in wide orbits are that there are likely to be several such planets for every planet that can give rise to a resonant event, and the probability of detection is not decreased in the same way ⁴, and can even be increased, by finite-source-size effects.

We will say that planetary orbits in which the separation between the star and planet is larger than roughly $1.5 R_E$ are “wide”. (See §2.3.) There are two types of events that provide evidence for the presence of planets in wide orbits. The first are isolated short-duration events due to the passage of the track of a distant star through the Einstein ring (or lensing region) of an intervening planet. The second are repeating events, in which the track of an individual lensed source passes through the lensing regions of more than one mass in the planetary system. Among repeating events, the probability is largest that the track of the source will pass through the lensing regions of the central star and one planet, but it can also pass through the rings of several planets. This paper focuses on isolated events; the companion paper (DiStefano & Scalzo 1998) concentrates on repeating events.

§2 lays the foundation for important parts of both papers. Although comprehensive introductions to microlensing can be found elsewhere (see, e.g., Paczyński 1996), §2 includes a brief general introduction to microlensing, as well as some material specific to the case of lensing by planets in wide orbits. Of particular importance are (1) the notion of the width of the lensing region, and (2) the role played by finite-source-size effects in the detection of planets in wide orbits. In §3 we address the question of whether planets in wide orbits are likely to exist and to serve as lenses. Event rates, and detectability issues are the subjects of §4. In §5 we discuss the detection strategies that can optimize the discovery of isolated short-duration events and test the hypothesis that they are due to lensing by planets; we focus on methods that involve modifications of existing observing programs. We sketch our conclusions in §6.

2. Technical Introduction

⁴The influence of finite source size on the detection of planets in the zone for resonant lensing depends on the location of the planet within the zone. The net effect has been calculated in most detail by Bennett & Rhie (1996). See DiStefano & Scalzo 1997, 1998 for a specific comparison of finite-source-size effects on the detection of planets in wide orbits with the detection of planets in the zone for resonant lensing.

2.1. Microlensing

Consider light traveling from a star located a distance D_S from us. If there is an intervening mass, M , located a distance D_L from us, gravity will deflect the light from the path it would otherwise have taken. The deflection leads to multiple and generally distorted images of the source. When D_S is on the order of tens of kpc and the lens is a solar-mass star, the maximum image separations are on the order of milliarcseconds, and are generally not resolvable. Changes in the amount of light received can be measured however, and these are also associated with lensing. The changes in flux are due to the motion of the source relative to the lens along the direction transverse to our line of sight. Let u represent the instantaneous distance between the source and the lens, as projected onto the lens plane. The instantaneous value of the magnification, A , is

$$A = \frac{u^2 + 2}{u\sqrt{u^2 + 4}}. \quad (1)$$

The magnification itself is due to the fact that, although the surface brightness of the source is not affected by gravitational lensing, the total surface area of the images does change as the projected position of the source moves closer to and farther away from the lens. In the above equation, u is measured in units of the Einstein radius, R_E .

$$R_E = \left[\frac{4 M G D_S x (1 - x)}{c^2} \right]^{\frac{1}{2}}, \quad (2)$$

where $x = D_L/D_S$. For $u = 1$, the magnification is equal to 1.34. This magnification is measurable; indeed, the Einstein radius is often used as a measure of the width of the lensing region. For a solar-mass lens, with $D_S = 10$ kpc and $x = 0.9$, $R_E = 2.7$ au.

In this paper and its companion (DiStefano & Scalzo 1998), we will often set D_S to 10 kpc and x to 0.9. This roughly corresponds to lensing by a star or planetary system in the Galactic Bulge of light from a more distant Bulge star.⁵ Because the greatest number of events have been discovered along the direction to the Bulge, and because many of these may in fact be events in which a Bulge star lenses light from a more distant Bulge star, these choices of the parameters may be the ones most relevant for the near-term discovery of planets via microlensing. The effects of other parameter choices can be gauged from Eq.

⁵Smaller distances to Bulge stars may apply to a large fraction of potential Bulge source stars. In these cases, x will typically be smaller, and the size of stellar and planetary Einstein rings will not differ substantially from the values we compute with $D_S = 10$ kpc and $x = 0.9$.

2. In general, making D_S larger, or choosing x closer to 0.5 (the value that maximizes the size of the Einstein ring for given choices of M and D_S), will increase the size of the Einstein radius, and will require planets to be located farther from the central star in order to be in “wide orbits” (§2.3).

A measure of the event duration is given by $(2 R_E)/v_T$, where v_T is the transverse velocity between source and lens. (See, e.g., Figure 6.) If $v_T = 100$ km/s, the event duration would be ~ 94 days, if the lens is of solar mass. If the lens is a Jupiter-mass (Earth-mass) planet, then the event duration would be ~ 3 days (4 hours). Clearly the short duration of events in which low-mass objects serve as lenses poses a challenge to the detection of planets via microlensing. We note, however, that when finite-source-size effects play a role, the relevant crossing time is more closely related to the time taken for the track of the source to move through a distance equal to the size of the source. If $R_S = 10 R_\odot$, and $v_T = 100$ km/s, $R_S/v_T \sim 19$ hours.

2.2. The Standard Strategy for Microlensing Detections of Planets

The strategy for planet discovery suggested by Mao & Paczyński (1991) and Gould & Loeb (1992) relies on observing an event due to lensing by the planetary system’s star, and detecting perturbations from the form given by Eq (1). Although the time scale for the stellar-lens event may be on the order of months, the perturbations due to planets are short-lived, lasting for times on the order of hours. The perturbations are due to the passage of the track of the source through or very near to the caustic structure associated with the presence of the planet. If the lensed source is point-like, then there is an infinite jump in the magnification as it passes through a caustic curve. Finite-source-size effects moderate the magnitude and characteristics of the change. Caustics are a feature associated with lens multiplicity. The case of a planet orbiting a star can be thought of as a binary system with an extreme mass ratio; generally we expect $q_p = m_p/m_*$ to be less than 10^{-2} or 10^{-3} . Because the mass of the planet is small, the caustic structures around the stellar position are small; this is why the perturbations of the stellar-lens light curve tend to be short-lived.

We will use the phrase “zone for resonant lensing” to refer to the region ($a_c < a < a_w$) in which caustic crossing events are most likely to occur. The spatial extent of the zone for resonant lensing was discussed by Paczyński (1996) and has been studied most recently by Wambsganss (1997). Our results on detecting planets in wide orbits do not rely heavily on the exact position of its inner boundary, a_w .

As presently implemented, the search for resonant planets is carried out in two steps. First, the monitoring teams (e.g., EROS, MACHO, OGLE) identify microlensing event candidates. They do this by monitoring the flux from $\mathcal{O}(10^7)$ stars regularly. Some of the monitored fields are not visited every night; others are the targets of regular nightly monitoring and may occasionally be re-visited even within a single night. If the monitoring teams discover that an otherwise non-variable star’s flux has increased significantly above baseline in several consecutive measurements, they typically issue an alert so that other observers can monitor the star more frequently. Follow-up teams have been formed to take systematic advantage of this opportunity (Udalski et al. 1994; Pratt et al. 1995; Albrow et al. 1996). Under favorable conditions, the follow-up teams carry out hourly monitoring with good photometry ($\sim 1\%$).

Although no promising candidates for events displaying clear signs of a planet lens have yet emerged from the data sets, there may be several reasons for this. For example, the detection efficiency may be low enough that a larger number of events need to be monitored in order to have a good chance of detecting a resonant event. It might also be that planets with the appropriate distance from the central star are not common. In addition, finite-source-size effects can wash out the signature of the short-time-scale perturbations characteristic of planetary mass ratios, decreasing the number of detectable events (Bennett & Rhie 1996).

2.3. Planets in Wide Orbits

The idea that the term “wide orbit” is meant to capture is that, as the orbital separation between the planet and the central star increases, caustic crossing events become less likely, while events in which a planet acts as an independent lens become more common. These latter are characteristic of planet-lens events, when the planet is in a wide orbit. If the track of the source passes through the Einstein ring of a single planet, we may discover an isolated event of short-duration. If the track of the lensed star passes through the Einstein rings of several masses in the planetary system, then we may hope to discover a repeating event. Although we have used the phrase “Einstein ring”, we note that, if the monitoring teams employ sensitive photometry, they can detect events in which the distance of closest approach to the lens is larger than an Einstein radius. It is therefore useful to introduce the notion of the “lensing region”: the region around the lens in the lens plane through which the source track must pass in order for a *detectable* event to occur.

We will say that a planet is in a “wide” orbit if its distance from the central star is large enough that the isomagnification contour associated with $A = 1.34$ is comprised of

two separated, closed curves, one centered near the star and the other centered near the planet. For mass ratios ranging from 10^{-5} to 10^{-3} , the critical orbital radius, a_w , beyond which an orbit becomes wide is roughly equal to $1.5R_E$. The choice of boundary, a_w , is of course somewhat arbitrary. Were the exact value of a_w important, then its dependence on the mass ratio would need to be taken into account. In fact, however, its exact value makes little difference to results on wide-orbit planets; these would be roughly the same had we chosen $a_w = 1.8 R_E$, or $a_w = 2.0 R_E$. The effect of increasing orbital separation near the boundaries of the zone for resonant lensing has been studied in detail by Wambsganss (1997), and readers are referred there for a detailed discussion and instructive graphics.

2.4. Width of the Lensing Region

Consider a lens of mass m_i . As mentioned above, the lensing region associated with this mass is that region in the lens plane through which the track of the source must pass in order for an *observable* event to occur. If the region for lensing by a particular star or planet is a disk, as it very nearly is when the planets are in wide orbits, the width, w_i , of the lensing region can be taken to be the radius of the lensing disk. Because the size of the lensing region relative to the Einstein ring is important, it is convenient to define

$$n_i = w_i / R_{E,i}. \quad (3)$$

n_i is the maximum impact parameter for an *observable* event, expressed in units of the Einstein radius. Note that n_i is not generally an integer. Large values of w_i facilitate planet detection in two ways. First, the event probability is proportional to w_i . Second, the larger the value of w_i , the longer typical events will be observable, and the better our chances of event detection.

The value of w_i can be made larger in two ways. The first is simply to increase the photometric sensitivity of the monitoring programs. Let A_{min} represent the minimum value of the peak magnification for which an event can be reliably detected. If $A_{min} = [1.58, 1.34, 1.06, 1.02]$, then $n = n_i = [0.76, 1, 2, 3]$.⁶ The general formula for n in terms of A_{min} , is

$$n = \sqrt{2} \left[\left(1 + \frac{1}{A_{min}^2 - 1} \right)^{\frac{1}{2}} - 1 \right]^{\frac{1}{2}}. \quad (4)$$

⁶Note that when finite-source-size effects are unimportant, n_i has the same value, n , for all lenses. Since, however, the role played by finite-source-size effects is specific to the lens, the value of n_i can be lens-specific.

The MACHO team has been driven to use a higher value of A_{min} (1.58) in order to eliminate the possibility that some forms of stellar variability may mistakenly be identified as microlensing (see, e.g., Purdue *et al.* 1995). As the star fields studied by the observing teams are monitored over longer times, however, unrecognizable stellar variability will become less of a potential problem, and the monitoring teams should be able to lower their values for A_{min} . The photometric sensitivity of the follow-up teams is such that, were they to attempt to identify new events in the fields where they are monitoring ongoing events, $A_{min} = 1.06$ ($w_i = 2 R_E$) could be achieved.

The second way in which w_i can be increased is if the size of the lensed star, as projected onto the lens plane, is comparable to or slightly larger than the size of the Einstein ring. Although, finite source size tends to decrease the value of the peak magnification, it can also increase the size of the lensing region, since some parts of the source may be near enough to the lens position to experience significant magnification, even when the center of the source is farther away than a point source would have to be in order to experience the same overall enhancement of flux.

2.5. Event Duration

Our primary interest is in the detection of isolated events of short-duration. Thus, our definition of event duration must be keyed to the amount of time during which there is a deviation from the baseline flux that would be discernible, given a particular level of photometric sensitivity. If events can be reliably detected when the peak magnification is A_{min} , then their start can certainly be detected before the magnification is A_{min} , and they can continue to be monitored after the magnification has fallen below A_{min} . In order to be conservative about the possibility of event discovery and detection, we will say that an event starts when the magnification rises above A_{min} , and ends when the magnification falls below A_{min} . Thus, the value of A_{min} determines the width, w_i , of the lensing region. w_i in turn plays a key role in setting the time, τ_i , during which an event is observable. We have $\tau_i = 2 \sqrt{w_i^2 - b^2} / v_T$, where b is the distance of closest approach. This definition of the event duration is directly related to the time during which the event can actually be monitored. It also makes it clear that any innovations that improve the photometric sensitivity of the observing programs may be important in increasing the observable duration of planet-lens events, thus increasing the chances that such events will be discovered and studied. When we present the results of our simulations, we will take $A_{min} = 1.06$.

2.6. Finite-Source-Size Effects

Because the Einstein radius scales as the square root of the lens mass, the Einstein radii of planet-mass lenses can be comparable to the projected radius of the source star. With $D_S = 10$ kpc and $x = 0.9$, $R_E = 17.8 R_\odot$ ($R_E = 1.0 R_\odot$) for a Jupiter-mass (Earth-mass) lens. When the size of the source becomes a significant fraction of the size of the Einstein ring, finite-source-size effects become important, and the magnification is not well described by the formula of Eq. 1. There are two important effects. First, the peak of the light curve becomes attenuated. The larger the source size relative to the Einstein ring, the more pronounced the attenuation. The maximum magnification possible for a disk of constant surface brightness, for example, is (Schneider, Ehlers, & Falco 1992)

$$A_{max} = \frac{\sqrt{R_S^2 + 4}}{R_S}. \quad (5)$$

This formula indicates that the size of the source must be fairly large in order for the peak magnification to be brought below the level of detectability, particularly if the photometric sensitivity is good. (Note that this applies to an isolated point mass; a more exact formula given by Gould & Gaucherel 1997.) For example, a magnification of $[1.34, 1.06, 1.02]$ can be achieved even if the size of the source is $\sim [2.2, 5.7, 10.] R_E$. Note that, when the lens is of Jupiter-mass (Earth-mass), then a star with radius roughly equal to $180 R_\odot$ ($10 R_\odot$) would fill a disk of radius $10 R_E$. (We have used $D_S = 10$ kpc and $x = 0.9$.) Thus, in spite of the decrease in peak magnification, the large majority of stars in the source population can, when are lensed by planets in wide orbits, produce events that should be detectable, at least with good photometry.

The second effect of finite source size on the light curve is to broaden the width of the perturbed region of the light curve, since some portions of the source may be significantly magnified well before the center of the source achieves its closest approach to the lens. The light curves can become almost flat-topped. Thus, magnifications close to the peak value may be sustained during the time it takes the source to travel several Einstein diameters, and the event appears to last longer than it would have had the lensed source been a point source. This can be a boon to the detectability of events due to planet-mass lenses, since one of the greatest barriers to detection is the short lifetime of the event.

Figure 1 displays a sequence of light curves for the case $R_S = 5 R_E$. When comparing the different light curves, note that the peak magnification does not change much, even as the distance of closest approach, D_{min} , changes from 0 to $4 R_E$. If the photometric sensitivity of the observations was such that events with peak magnification above 1.06 could be detected, then all of these events would be detected. The primary difference as




Fig. 1.— In each panel, the projected size of the source in the lens plane is $5 R_{E,i}$. The distance of closest approach, $D_{\min}$, decreases from $4 R_{E,i}$ in the upper panel to the 0 in the lower panel. Note that the peak magnification is larger than 1.06 in all cases. Furthermore, if the measured event duration is the time during which the magnification is larger than 1.06, then these events last up to 2.5 (bottom panel) times as long as they would have, had the lensed source been a point source. We have assumed that the disk of the lensed source has constant surface brightness.




Fig. 2.— The source radius, R_S , is plotted against D_{min} , the distance of closest approach. Both quantities are expressed in units of the Einstein ring radius. Each point corresponds to an event in which the peak magnification was roughly equal to the value listed on the upper right of each curve. For each value of R_S , only the points with the largest values of D_{min} leading to the listed magnification are shown. We have assumed that the disk of the lensed source has constant surface brightness. The value of D_{min} for a point source is shown, for $A_{min} = 1.02, 1.06$, and 2.0 [vertical lines].

D_{min} increases, is that the time duration of the observed event decreases. Even so, for D_{min} as large as $4 R_E$, the observed duration is still roughly equal to the time a point-source light curve would remain above $A = 1.06$. The relationship between D_{min} and R_S is explored more systematically in Figure 2. The calculations used to produce these figures assumed that the surface brightness of the lensed star was constant across the face of the star. More realistic stellar profiles will change the details, but the general features are robust. The implications are that: (1) depending on the distribution of source sizes, the detection probabilities can be significantly increased by finite-source-size effects, and (2) events that can be detected can also generally be monitored for longer periods of time.

Gould (1994) and Peng (1997) were able to find point-source fits to light curves associated with extended sources. This degeneracy of the light curve shape is not expected to be problematic, however, when the source is larger relative to the Einstein ring, as it will be in many of the cases relevant to the discovery of wide-orbit planets (DiStefano 1998a). Moreover, even if there are light curve degeneracies, these will not necessarily prevent the identification of individual events as having been influenced by finite-source-size effects, since spectral information can break such degeneracies. Real stellar disks exhibit brightness profiles and spectra that have spatial structure. When a star for which $r = R_S/R_{E,i}$ is not negligible is lensed, the observed spectrum, and therefore the light curve as observed at different wavelengths, will be time-dependent. This effect can be used to study the lensed star. (See, e.g., Witt & Mao 1994, Witt 1995, Gould 1994, Loeb & Sasselov 1995, Simmons, Willis & Newsam 1995, Gould & Welch 1996, Sasselov 1997, Heyrovsky & Loeb 1997, and, for an observation, Alcock *et al.* 1997.) This time dependence also helps to confirm the interpretation of the event as due to microlensing of a source whose size has influenced the shape of the light curve. Indeed, when the source is large relative to the Einstein ring, it is as if a magnifying glass were scanning the face of the lensed star, thus providing a good deal of information about the source star and about the event. Spectral studies, even during short-duration events can therefore be valuable in breaking the degeneracy for individual events, as can multi-color photometry. Even if spectral information is not available, however, statistical analysis of the ensemble of events would be able to indicate that finite-source-size effects had played a role. In particular, the relationship between peak magnitude and duration would be different than it would be for a set of true Paczyński light curves; for large sources, the peak magnification is closely related to source size, while the duration continues to have a more direct relationship to the distance of closest approach.

2.7. Resonant, Repeating, Overlap, and Isolated Events

In the companion paper (DiStefano & Scalzo 1998) we provide an overview of how planet-lens events of different types (resonant, wide repeating events, wide isolated events) complement each other. To orient the reader, we provide a thumbnail sketch here.

Orbital separation: When the separation, a , between the central star and a planet is much smaller than the star’s Einstein radius (typically $a < 0.5 R_E$), the large majority of events in which the planetary system serves as a lens will be indistinguishable from stellar-lens events in which an isolated star serves as a lens. As a increases, there is a marked increase in the fraction of stellar-lens events displaying distinctive “resonant” perturbations linked to the presence of the planet. As a continues to increase, however, such events again become rare: i.e., more distant planets are not in the zone for resonant lensing.

For planets located just beyond the zone for resonant lensing, the probability of repeating events achieves its maximum value. In fact, if the photometric sensitivity is good, some repeating events (in which the source passes through the lensing regions of both the star and a planet), will exhibit light curves in which the magnification has not yet returned to baseline levels when the influence of a second lens becomes apparent. We refer to this subset of repeating events as “overlap” events.

Conceptually, overlap events are most closely related to repeating events, since they occur when the lensing regions of two lenses are transited by a distant source. Phenomenologically, however, they are “isolated” events, in that the associated light curves exhibit only a single excursion from the baseline flux. If the photometry is sensitive enough, there is no wait-time between events, and this increases the probability they will be identified, relative to the probability of detecting other repeating events. Because they are relatively long-lived, with their time duration keyed to the duration of stellar-lens events, they are also more likely to be detected than isolated events of short-duration. In this paper we focus primarily on isolated events of short-duration, since these will be most challenging to detect. Because they appear as isolated events, however, we do include overlap events in the computations described and summarized in §4.

As a increases further, all repeating events display genuine repeats; i.e., they restart after a fall back to baseline following an initial event. The probability of repeating events, which scales as $1/a$, decreases, and the probability of isolated events of short-duration increases. At large separations, isolated events of short-duration dominate.

Thus, each type of event probes a different region of planetary systems that act as lenses. (We note, however, that because different spatial orientations are expected, the regions we study with each type of event can intersect the regions explored by the others.) Detections

of each type of event therefore complement detections of the others, with the ensemble of all events providing information about the spatial structures of and mass distributions within planetary systems. In addition, discoveries of resonant and wide events are subject to different selection effects, allowing them to complement each other in another way.

The light curves: The light curves associated with resonant events are essentially stellar-lens light curves, exhibiting short-lived perturbations due to the presence of a planet. In principle, the mass ratio between the planet and star, and the projected value of their orbital separation, can be derived from the light curve; in practice, there are degeneracies in the physical solutions that must be considered (Gaudi 1997, Gaudi & Gould 1997). The light curves associated with wide-orbit events exhibit at least one peak that is of relatively short duration, since at least one lens is a planet and not a star. For repeating events, as for resonant events, the mass ratio and orbital separation can be constrained based on the light curve alone. As always, there are degeneracies, but the fact that we see two separate encounters in which one source is lensed may help to resolve those degeneracies involving blending and finite source size. For isolated events, the light curve alone generally provides no specific information about the orbital separation, and we can only derive a lower limit on the distance between the central star and the planet, expressed in terms of the size of the star’s Einstein radius. If, however, the spectral type of the central star can be determined (which may be possible if it contributes a significant fraction of the baseline flux), and/or if finite-source-size effects are important, then we can learn more even about planet lenses giving rise to isolated events. (See the discussion below.) In addition, for some isolated events, the form of the light curve will deviate from the standard form in a measurable way. In a very small fraction of cases there may be caustic crossings, with the light curve form also significantly influenced by finite-source-size effects. In others, the light curve will be well-fit by the formulae of DiStefano & Mao, allowing the projected orbital separation to be derived.

Novel aspects of planet detection via microlensing: In addition to the wide-orbit channel for detection, there are other aspects of planet detection via microlensing that have been little-explored. One of these has to do with follow-up work that can be done for planet-lens events, to learn more about the planetary system that served as a lens. While it has widely been assumed that little can be learned about planetary systems discovered through lensing, the combination of blending and finite-source-size effects can play a powerful role in helping us to determine the spectral type of the central star and the masses of any planets involved in the lensing event. We touch on this here (and in DiStefano & Scalzo 1997, 1998), but discuss the possibilities more fully in DiStefano 1998b. The relevance to this paper is that it is possible that observations designed to complement light curve studies can help us to learn more about isolated events of short-duration. The

additional information can, under favorable circumstances, include the spectral type of the central star and the mass of the planet lens. Another interesting question is whether the planets discovered via microlensing are likely to harbor life; this is discussed in DiStefano (1998a). Finally, we note that it may be possible to detect fine structure within the lensing planetary system, in the form of moons orbiting planets and asteroid or cometary belts (DiStefano & Keeton 1998).⁷

3. Do We Expect To Find Planets in Wide Orbits?

The excitement about resonant lensing by planets was fueled, at least in part, by a wonderful coincidence. Gould & Loeb (1992) noted that, if a system identical to our Solar System happened to be located halfway between our position and the center of the Bulge, and if the system were viewed face-on, the separation of the planet corresponding to Jupiter from the system’s star would be very close to the value of R_E associated with the mass of the star. That is, Jupiter would be in the zone for resonant lensing. There are, however, two reasons to be cautious about using this example to limit the search for planets to those in resonant orbits. First, we do not know that the spatial relationship between the Sun and Jupiter is an example of a universal property of planetary systems. Mindful of this, Bennett & Rhie (1996) have constructed a “power-of-2” planetary model, in which the separation between each planet and the central star increases by a factor of two for each successive planet. In such a model, most planetary systems can be expected to contain one planet in a resonant orbit. Each such planetary system contains several ($\mathcal{O}(10)$) planets in wide orbits. A second reason to avoid limiting the microlensing searches to resonant planets is that the value of the Einstein radius depends not only on the stellar mass, but also on the relative positions of both source and lens to the observer.

In Figure 3 we consider planets with orbital separation $a = 2a_w = 3R_E$. For these systems we show the relationship between the stellar mass, M , and x , when the orbital period is fixed. We have assumed that the orbital plane is the same as the lens plane, and show those values of x for which both lens and lensed source are located in the source galaxy (the Bulge, the Magellanic Clouds, or M31). If the stellar mass is in the range from 0.1 to $10 M_\odot$, the orbital periods of planets in wide orbits range from a few years to a few hundred years, with larger values more typical for more distant galaxies.

⁷All of these topics, together with both isolated and repeating wide-orbit events, are covered in DiStefano & Scalzo 1997 as well. The referee of that paper suggested that it would be preferable to discuss the separate ideas in separate papers, thus leading to the sequence of papers described here. The collection of individual papers includes some new work.




Fig. 3.— Each curve corresponds to a fixed value of the orbital period; shown are curves for 1 yr, 10 yrs, 100 yrs, 1000 yrs. Each curve illustrates the relationship between the stellar mass, M , and x for a planet in a wide orbit, with $a = 3 R_E$. In each panel the range of the variable x corresponds to lenses located within the same galaxy (Bulge, Magellanic Clouds, M31 [left to right]) as the lensed stars. Note that the range of masses and orbital periods best explored via wide lensing events is different for different source galaxies.

Table 1. Planets in wide and resonant orbits in known and model systems;
 $D_s = 10$ kpc, $x = 0.9$.

Planetary system	Planet	a ⁽¹⁾	$\%_{close}$ ⁽²⁾	$\%_{res}$ ⁽³⁾	$\%_{wide}$ ⁽⁴⁾
Solar system:	Jupiter	5.2	5.9	21.0	73.1
	Saturn	9.5	1.7	4.6	93.7
	Uranus	19.2	0.4	1.0	98.6
	Neptune	30.1	0.2	0.4	99.4
	Pluto	39.8	0.1	0.2	99.7
Known extrasolar systems:	55 Cnc	4.0	11.0	89.0	0.0
	HD 29587	2.5	37.0	63.0	0.0
	PSR B1620-26	38	0.1	0.3	99.6
	Gl 229b	40	0.1	0.2	99.7
Theoretical power-of-2 system: ⁽⁵⁾	1	4.8	7.0	27.0	66.0
	2	9.6	1.6	4.5	93.9
	3	19.2	0.4	1.0	98.6
Theoretical power-of-3 system: ⁽⁶⁾	1	4.8	7.0	27.0	66.0
	2	14.4	0.7	1.9	97.4
	3	43.2	0.1	0.2	99.7

⁽¹⁾Separation of the planet from the central star in astronomical units (AU). We consider circular orbits. See the text for the criteria used to select the planets listed here.

⁽²⁾Percentage of the time during which the planet is “close” ($a < 0.8R_E$), averaged over all inclinations.

⁽³⁾Percentage of the time during which the planet is “resonant” ($0.8R_E < a < 1.5R_E$), averaged over all inclinations.

⁽⁴⁾Percentage of the time during which the planet is “wide” ($a > 1.5R_E$), averaged over all inclinations. In the power-of-2 and power-of-3 models, all planets beyond the third had a probability greater than 99% of being located more than $1.5R_E$ from the central star.

⁽⁵⁾The orbital separation increases by a factor of 2 for each successive planet. When we averaged over possible positions for the innermost planet (from 0.3 AU to 2.0 AU), keeping the number of planets constant at 12, we found that 1.90 planets on average were close; 0.925 planets on average were resonant; and 9.17 planets on average were wide.

⁽⁶⁾ The orbital separation increases by a factor of 3 for each successive planet. When we averaged over possible positions for the innermost planet (from 0.3 AU to 2.0 AU), keeping the number of planets constant at 7, we found that 1.41 planets on average were close; 0.535 planets on average were resonant; and 5.06 planets on average were wide.

In addition, the projected value of the separation depends on the angle, α , between the normal to the orbital plane and our line of sight. Consider an orbit with $a > a_w$. Let α represent the angle between the normal to the plane of the orbit and the normal to the lens plane. For $\alpha < \alpha_{min}$, where

$$\alpha_{min} = \cos^{-1}(a_w/a), \quad (6)$$

the projected value of the orbital separation is always greater than a_w . If $\alpha > \alpha_{min}$, then for a circular orbit, the fraction of time for which the projected separation is greater than a_w (so that the planet can be viewed as wide) is

$$f = \frac{2}{\pi} \cos^{-1} \left(\frac{\sqrt{\cos^2 \alpha_{min} - \cos^2 \alpha}}{\sin \alpha} \right). \quad (7)$$

If the orbit is eccentric, then the separation between the planet and star will be wide for an even greater fraction of the time.

Thus, even if planetary systems had uniform properties, the fact that they are located at different spatial positions and are tilted at different angles relative to our line of sight would still not favor resonant orbits over others. Bennett & Rhie’s power-of-2 model, or something like it, is necessary to ensure that most planetary systems have one planet in the zone ($a_c < a < a_w$) associated with resonant lensing. This is because $a_w/a_c \approx 2$; thus, if the i^{th} planet has separation a_i from the central star, and the power-of-2 system is inclined so that $a_i \cos \alpha \approx a_c$, we have $a_{i+1} \cos \alpha \approx a_w$.

The known planetary and brown-dwarf binary systems are considered in Table 1, together with a “power of 2” model and a “power of 3” model. The known systems included in the table were selected from the Encyclopedia of Extrasolar Planets (accessible from <http://www.obspm.fr/planets>) Two criteria were used: (1) if the planetary system were placed in the Bulge and viewed face-on, the planet (or brown dwarf) listed had to be in an orbit that would either be wide or located in the zone for resonant lensing, (2) the existence of the planet or brown dwarf needed to be listed as “confirmed”. In the power-of-2 and power-of-3 model, the first planet listed is the innermost planet that would be in a wide orbit, were the planetary system to be placed in the Bulge and viewed face-on. We note that the next planet inward would have a good chance of being viewed in the zone for resonant lensing. Indeed for both of these theoretical models, there is some ambiguity associated with the arbitrarily-chosen position of the closest planet to the central star. Because of this, we carried out a set of calculations in which we averaged over the position of the innermost planet. We found that, if we truncate the radius of the planetary system at $\sim 10^{17}$ cm, there are on average $\sim 9 - 10$ planets in wide orbits for every planet located in the zone for resonant lensing. To derive the numbers shown in the table, we averaged over inclination

angle α , and used equations (6) and (7) to determine the fraction of time any given planet would be in a wide orbit. As above, we have assumed that the orbits are circular.

The considerations in this section show that, under a range of reasonable assumptions, planets in wide orbits are very likely to exist. In fact the number of planets in wide orbits may be as much as an order of magnitude larger than the number of planets in resonant orbits. The relative contribution of these two classes of planet lenses to the detection rate depends on our ability to identify the associated events.

4. The Rate of Detectable Isolated Planet-Lens Events

4.1. Normalization of Event Rates

We will be interested in the rate at which detectable events occur, and how that rate is influenced by changes in the detection strategy. The Einstein radius of the central star provides a convenient normalization. The rate of encounters in which the central star serves as a lens, with the magnification due to the central star achieving a peak value of at least 1.34 ($A_{min} = 1.34$), is proportional to $2 R_E$. Because the width of the central star's lensing region (when $A_{min} = 1.34$) will play an important role, we define $w_{0,0}$ to be equal to R_E . If the photometric sensitivity is such that A_{min} can be smaller than 1.34 (i.e., $w_i > R_E$, $n > 1$), so that the effective width of events involving the central star is larger ($w_0 > w_{0,0}$), the rate of all detected events, including those due to the central star, will increase. Nevertheless, $2w_{0,0}$ is a convenient normalization constant, because it allows us to compare the rates we compute to the presently-measured rate of events. This is as follows. Along the direction to the Bulge, events are being discovered by the MACHO team at a rate of roughly 50 per year. The present detection criteria used by the MACHO team are strict: $A_{min} = 1.58$, corresponding to n (in Eq. 3) being set equal to ~ 0.76 . In addition, other teams, surveying different fields with somewhat different strategies, are also surveying the Bulge. For example, the OGLE team, which in its first incarnation discovered ~ 12 Bulge events, has recently brought a new telescope on-line (Udalski, Kubiak & Szymanski 1997; Paczyński *et al.* 1997). It is therefore reasonable to assume 75 – 100 events of the type we use for our normalization per year along the direction to the Bulge. Thus, because we use $2w_{0,0}$ to normalize our results, when we find that a particular detection strategy leads to a rate of detectable events of a certain type (e.g., events with 2 repetitions) equal to $p\%$, this means that between $0.75p$ and p such events could be discovered per year along the direction to the Bulge.

4.2. The Rate of Isolated Events

When a wide planetary system serves as a lens, the most common type of event is one in which the track of the source passes through the Einstein ring of a single mass. Because the rate of encounters associated with a given lens scales as the square root of its mass, encounters in which the star serves as a lens are the most frequent. Let P_1 represent the rate of isolated short-duration events due to planet lenses. P_1 is proportional to the linear dimensions of the lensing region.

$$P_1 \sim 2 \sum_{i=1}^N w_i \sim 2 \sum_{i=1}^N n_i R_{E,i} \sim \sum_{i=1}^N n_i \sqrt{q_i}, \quad (8)$$

where the sum is over all of the system's planets, and $q_i = m_i/m_*$. When finite-source-size effects are not important, $w_i = n R_{E,i}$. The normalization described in §4.1, yields a normalized rate of isolated planet-lens events:

$$P_1 = \sum_{i=1}^N n_i \sqrt{q_i} = n \sum_{i=1}^N \sqrt{q_i}, \quad (9)$$

Note that, because the event rate scales with n , improvements in photometric sensitivity play an important role in optimizing the rate of detectable events, even when finite-source-size effects are not important. When finite-source-size effects do play a role, the increase in rate can be even larger for some of the terms on the right hand side of Equation 8.

The expression for P_1 given in Eq. 9 overcounts isolated events slightly, because the rate of repeating events must be subtracted from it. Because, however, isolated events involving a specific lens are generally 1 – 2 orders of magnitude more common than repeating events involving the same lens, (DiStefano & Mao 1996; DiStefano & Scalzo 1997, 1998), P_1 generally provides a good estimate of the rate of isolated events involving planets.

The time duration of isolated planetary-lens events, relative to the time duration of stellar-lens events, also scales as $\sqrt{q_i}$. The implication is that a distribution of events due to lensing by stars with wide-orbit planets is necessarily accompanied by a distribution of shorter-duration events. The fraction of events in the latter distribution is proportional to the average value of $\sqrt{q}$, and the position of the peak or peaks also provides a measure of the mass ratios typical of planetary systems.

Possible forms of the distribution of event durations are illustrated in Figures 4 and 5. (The rates corresponding to the integrated area under the curves for planet-lens events are discussed in §4.3.) These figures show results from a set of Monte-Carlo simulations in which 3 types of planetary systems were placed in the Bulge and served as lenses for more distant Bulge stars ($D_S = 10$ kpc, $x = 0.9$). The details of the simulations and a

more complete description of the results are provided in the companion paper (DiStefano & Scalzo 1998). Shown are the results derived when the planetary system lens was (a) identical to the Solar System (top panel), (b) a power-of-3 system (middle panel), and (c) a power-of-2 system (bottom panel). (Note that, in the power-of- n models, the orbital separation between planets and the central star increases by a factor of n proceeding outward from the central star.) Each planetary system was considered as a lens in numerical experiments that generated thousands of random orientations of the orbital plane with respect to the lens plane, and for each, considered the passage of source tracks coming in at random angles, with velocities chosen from a Gaussian distribution centered at 150 km/s, with width equal to 50 km/s.

We show the (unnormalized) distribution of event durations, defining the duration to be the time during which the magnification was above $A_{min} = 1.06$. In all cases, a long-duration peak in the distribution, due to lensing by the central star, is necessarily accompanied by one or more smaller peaks due to short-duration planet-lens events. The difference between Figure 4 and Figure 5 is that different sets of criteria were used to determine whether an event is detectable. Criteria A (Figure 4) require the light curve to exhibit magnification greater than $A = 1.34$ (source-lens separation less than $1.0R_E$) for at least 1 day in order for the lensing event to be detected. Criteria C (Figure 5) require only that the magnification be greater than 1.06 (source-lens separation less than $2.0R_E$), and no minimum event duration is required. Criteria C are more ambitious, and allow the effects of lensing by several planets to be seen. In the case of the Solar System, there are peaks due to lensing by each of the outer planets, although some of the structure is blurred as the peaks run together. It is interesting to note that most of the peak structure actually comes from Saturn and the planets beyond it, rather than from Jupiter. The reason for this is that the separation between Jupiter and the sun is so close to a_w that, for some orientations of the orbital plane with respect to the lens plane, Jupiter was in the zone for resonant lensing at least part of the time. Furthermore, even when it was in a wide orbit, its lensing region sometimes overlapped that of the Sun. That is, before the magnification could slip below 1.06 after Jupiter served as a lens, the track of the source would already be in the lensing region of the Sun (or *vice versa*). The result would be one long event with the light curve exhibiting two peaks. The dotted lines correspond to such “overlap” events. Although the source track passes through the lensing regions of two lenses, there is just one continuous deviation from baseline. Overlap events appear to be isolated events with unusual morphologies that last for times comparable to the durations of stellar-lens events. Overlap events are discussed in the companion paper (DiStefano & Scalzo 1998). Like isolated events of short-duration, overlap events are necessary adjuncts of the presence of wide-orbit planets. They are a feature of the power-of- n simulations as well, when there

was often an overlap between the lensing region of the central star and the innermost wide planet.

In the power-of-2 and power-of-3 models, all of the planets were assumed to be of Jupiter mass, so there is just one statistically significant peak due to planet lenses; the area under this peak is large, however, because it includes the contributions from all of the systems’ wide planets.

4.3. Rates of Events Associated with Known Systems

The planetary system we know best is our own. Not only do we know that the Sun is accompanied by a planet, but we know that there is a set of planets distributed over several tens of astronomical units (au). If a planetary system identical to the Solar System is located in the Bulge, the Magellanic Clouds, or in M31, several of its planets (generally Jupiter and all of the outer planets) are in wide orbits most of the time, for most orientations of the ecliptic on the sky. If a lensing event with a solar mass lens lasts for 100 days, then an event in which [Jupiter, Saturn, Uranus, Neptune, Pluto] serves as a lens lasts for [3.08, 1.68, 0.66, 0.71, 0.01] days. If all stars were as massive as the sun, and were accompanied by a planetary system identical to ours, then the durations given in the previous sentence could be translated directly into the percentage of events in which each of the planets in turn could be expected to serve as a lens. The number of planet-lens events per year would be roughly 2 – 4 in Baade’s window (for $A_{min} = 1.34$), and $\sim 5 - 10$ (for $A_{min} = 1.06$). With the present monitoring strategy, events lasting 1 day or less would not be subjected to detailed enough monitoring to unambiguously identify them as microlensing events.

To make contact with specific detection criteria, we refer to the Monte-Carlo calculations corresponding to Figures 4 and 5. These yield a rate of isolated events of only 0.3 per year for the Solar System, when detection criteria A are used. This is because The lensing region of Jupiter actually overlaps that of the Sun for most orientations of the orbital plane on the sky. Thus, Jupiter is detected through overlap events, which occur at a rate of 2.2 per year when detection criteria A are employed. The only other outer planet that can produce an event lasting longer than a day is Saturn, and not every event due to Saturn lasts this long. Most of the overlap events should be detectable, and should also provide some evidence of the presence of Jupiter. The set of detection criteria C leads to 3.1 short-duration isolated events per year and 4.2 overlap events. Thus, over a period of 10 years, dozens of planet-lens events could be detected.




Fig. 4.— Distributions of durations for isolated (non-repeating) events using the detection criteria of set A (the light curve must exhibit magnification greater than $A = 1.34$ [source-lens separation less than $1.0R_E$] for at least 1 day in order for the lensing event to be detected). In each panel, the peak on the left is due to isolated planet-lens events, and the peak on the right is due to the central star. Overlap events, produced almost exclusively by encounters involving the innermost wide planet and the central star, are shown by the dotted lines. (See DiStefano & Scalzo 1998.) The truncation of the large-duration end of the overlap distributions in the power-of-3 and power-of-2 models is due solely to the poorer statistics achieved in the simulations of these models; all of the distributions were scaled to the linear sampling achieved for the Solar System. In the Solar System (power-of-3, power-of-2) model (top panel [middle panel, bottom panel]), there were 0.3 (13.0, 23.7) isolated events of short-duration per year and 2.1 (1.9, 2.2) overlap events.




Fig. 5.— Distributions of durations for isolated (non-repeating) events using the detection criteria of set C (the source-lens separation must be smaller than $2.0R_E$, and no minimum event duration is required). Here, because an encounter did not have to have a minimum duration in order to be detected, the peaks tend to have low-duration tails which can run together. The right-most peak in each panel is due to the central star of the system, with overlap events involving the innermost wide planet being shown by dotted lines (DiStefano & Scalzo 1998); the peaks on the left are due to planetary encounters. The central feature in the top panel is a superposition of peaks corresponding to Uranus, Neptune, and Saturn; the encounters due to Saturn are visible as a shoulder at about 1.6 days. In the Solar System (power-of-3, power-of-2) model (top panel [middle panel, bottom panel]), there were 3.1 (30.4, 55.9) isolated events of short-duration per year and 4.2 (3.6, 4.0) overlap events.

The event durations and rates scale with w_i , and w_i depends on finite-source-size effects. We therefore note that, if the Einstein radius of an Earth-mass planet is $\sim 1.0 R_\odot$, then the Einstein radius of [Jupiter, Saturn, Uranus, Neptune, Pluto] is [17.8, 9.8, 4.1, 3.8, 0.04] $R_\odot$. Pluto would very likely not be discovered via microlensing. Jupiter, Saturn, Uranus, Neptune would be detectable. Furthermore, were the lensed star to be massive and/or evolving, the time duration of the events could be significantly extended by finite-source-size effects. Thus, depending on the distribution of radii among the source population, the rate of detectable events could be significantly higher than given above.

We now know of planetary systems in addition to our own Solar System. Among other systems in which the presence of a brown dwarf or planet-mass is confirmed, there are two in which the brown dwarf or planet is in an orbit that would be wide, were the system to be located in the Bulge and serve as a lens for a more distant Bulge star ($D_S = 10$ kpc, $x = 0.9$). These are Gl 229 (a brown dwarf system) and PSR B1620-26. If all stars had similar companions to the brown dwarf in Gl 229 (the planet in PSR B1620-26), then there would be ~ 20 (10) isolated short-duration events per year along the direction to the Bulge (for $A_{min} = 1.34$,) and ~ 40 (20) isolated short-duration events per year (for $A_{min} = 1.06$).

It is simply not known whether our planetary system or any of the others which we have now begun to study are typical of planetary systems elsewhere in our Galaxy or in other galaxies. It may therefore be instructive to consider theoretical constructs. In the power-of- n model, we assume that there is a planet of mass m located within 1 au ($a_0 < 1$ au) of the central star, and other planets located at orbits of radii $n^i a_0$ for i ranging from 1 to the maximum value consistent with a bound orbit. For the power-of-2, 3, 4 model, with $A_{min} = 1.34$ (1.06), there would be $\sim 26, 15, 13 \sqrt{m/m_J}$ (52, 30, 26 $\sqrt{m/m_J}$) isolated short-duration events per year toward the Bulge per year. The results of Monte Carlo simulations are consistent with these predictions; for the power-of-2 and power-of-3 models the rates are given in the captions to Figures 4 and 5.

The common factor in all these results is that the event rates are large enough that, if planetary systems are a common phenomenon, we should be able to see evidence of planetary systems in the distributions of event durations.

4.4. The Detection and Identification of Isolated Planet-Lens Events

Are the observing teams capable of discovering events of such short durations that the lens could be a planet? If, so, and if planet-mass objects do serve as lenses, will we be able to identify the associated peak of short-duration events in the distribution of event

durations? This would allow us to minimize contamination of the planet-lens signal due to the short-duration tail associated with stellar lenses. Finally, and perhaps most interesting, can some individual isolated events be reliably identified as planet-lens events? If so, then we can eliminate some of the possible contamination due to lensing by low-mass MACHOs.

4.4.1. *Observations of Short-Duration Events*

The shortest event on record lasted approximately 2 days. To better understand what is known so far, we have considered events along the direction to the Bulge, since the MACHO team has already accumulated a store of over 150 Bulge events about which some information is publicly available. Specifically, we have used their paper on 45 Bulge events monitored in 1993 (Alcock *et al.* 1997a), and their “alert” web pages (<http://darkstar.astro.washington.edu>), which list approximate durations for many of the 136 Bulge events observed during 1995, 1996, and 1997. Altogether we found (up to October 1997) 148 events (a) which are apparently due to lensing by a point-mass, and (b) for which approximate durations are available. (We note however, that the durations posted on the web site come from fits that may be refined in the future.) The distribution of event durations is plotted in Figure 6. Note the appearance of the short-duration peak.

In light of Figure 6 we may ask whether the MACHO team has already begun to discover concrete evidence for a signature due to planets in the Bulge. This may be the case. (In fact Bennett *et al.* [1996a] have conjectured that one short event may indeed be due to a planet in a wide orbit.) We emphasize, however, that a good deal of further work would be required to clearly establish it (§4.6, DiStefano & Scalzo 1997, 1998, DiStefano 1998a,b). One certain conclusion we can draw from the MACHO Bulge data is that the team has proved that it has the ability to detect short-duration events. This is encouraging, since the detection strategies presently used are not optimized for the discovery of short time-scale events.

4.4.2. *Identifying Peaks In The Distribution of Short Events*

Consider the distribution of event durations. Is it likely that a peak due to the presence of lenses with small masses will be hidden by or confused with the short-duration tail of encounters due to lensing by stars?

Three circumstances can make a stellar-lens encounter have a short measured duration.

- (1) The relative velocity v_T an observer would measure between the source and the




Fig. 6.— Distribution of event durations for the MACHO team’s set of Bulge events. Included is data from 1993 (Alcock *et al* 1997a), and the “alert” web page data for events which occurred in 1995, 1996, and 1997. All 148 events which were not listed as binary events, and for which estimates of event durations had been, made are included. Note that the definition of duration used here is $(2 R_E/v_T)$.

lens may be exceptionally large. Since, however, the distribution of transverse velocities for planet-lens and stellar-lens encounters should be roughly the same, we should be able to disentangle velocity effects, at least for statistical samples of encounters.

(2) The track of the source may just graze the lens’ Einstein ring. In this case, the peak value of the magnification allows us to measure the distance of closest approach, D_{min} , and to determine that, although short, the observed encounter was nevertheless due to the presence of a lens with a larger Einstein ring than expected for a planet. These corrections are implemented by the observing teams.

(3) Light from the lensed source could be blended with light from other sources along the line of sight. If, however, the effects of blending are dramatic enough to shorten the duration of an event significantly, then we should be able to detect evidence for the blending in other ways, and to subtract its effects from the event before comparing its duration with that of other events. Because blending can provide a valuable tool for the study of planet-lenses, we discuss it in more detail in separate papers (DiStefano1998a,b).

We conclude that, if the present generation of monitoring programs discover a set of isolated planet-lens events, they should be able to reliably separate the signal from the contribution due to stellar lenses. As we see from Figure 2, the small short-duration “peak” in the MACHO team’s real data set is distinguishable from the rest of the distribution of event durations.

4.4.3. *Is an Isolated Event Really Due to Lensing by a Planet?*

Numerical simulations (Figures 4 and 5), the arguments above, and the data from the MACHO team, all indicate that the observing teams are likely to be able to find convincing evidence of short-duration events, if there is a population of low-mass lenses. Will the teams be able to determine, however, that the events are due to planets, rather than to low-mass MACHOs? A related question is whether a signal due to low-mass MACHOs could be obscured due to lensing by ordinary planetary systems.⁸ It is difficult to come up with comprehensive answers to such questions, because the distinction between planets and MACHOs is not always clear. For example, when low-mass objects are discovered to be orbiting a pulsar, they have been called “planets”. On the other hand, non-luminous, stellar

⁸Had a large portion of the Galactic Halo had been in the form of planet-mass compact objects, then the planet signal certainly could not obscure the MACHO signal. The data collected to date, however, seem to rule this possibility out.

remnants in the Galactic Halo and any planet-mass companions they may have would likely be considered as MACHOs. Single, non-luminous planet-mass objects may be referred to as MACHOs, but some of these may have been ejected from planetary systems with luminous central stars.

The most stringent criterion we can use to classify a low-mass object as a planet is that it should be in orbit with a luminous star. This definition actually gives us a way to test whether a lensing event was due to lensing by a planet.

If the central star of a planetary system lens is luminous, light from the lensed star will be blended with it. Let us suppose that we observe a short-duration isolated event, and that we can establish the following facts about it.

- (1) Not all of the flux emanating from the line of sight to the event is emitted by the lensed star.
- (2) At least some of the blending is due to light from a star whose position cannot be resolved from that of the lensed star, even with sub-arcsecond resolution.

In such a case, given the stellar surface density in the Bulge and LMC, it is almost certain that the additional baseline flux emanates from the lens system. This would definitively establish that the lens giving rise to the short-duration event was a planet. In fact, if the amount of light emanating from the planetary system’s central star can be quantified as a function of wavelength, we may be able to determine its spectral type. The observability of blended events is discussed in DiStefano 1998a,b. We find that the majority of events should still be detectable even when there is blending.

A test of the mass range of the lens (but not the planet-lens hypothesis) could be provided by finite-source-size effects. When spectroscopic studies of the lensed source can establish its real physical size, finite-source-size effects may then allow us to derive the true size of the Einstein ring. This means that, if the value of r , the ratio between the radius of the lensed star and the Einstein radius of the lens, can be measured from the data, v_T can be determined. Furthermore the degeneracy in the lens mass associated with the Paczyński light curve is partially broken, since knowing the value of R_E , and, through spectral studies, the value of D_S , allows us to infer

$$M x (1 - x) = C, \tag{10}$$

where C is a constant whose value is measured. Given the distribution of likely values of x (from 0.9 to 0.99, for example, in the Bulge) this generally constrains M to within a factor of ~ 5 . Thus, the hypothesis that any given event is due to lensing by a planetary-mass can be meaningfully tested. Furthermore, if the spectral type of the central star of the lens

system has been determined through the study of blending effects, then x can be completely determined.

5. Detection Strategies

Because good photometric sensitivity increases the rate of observable events and lengthens the duration of events that are observed, the the bottom line is that good photometry is the key to increasing the detection rate of planets in wide orbits. Monitoring that is consistently more frequent than that presently carried out by the monitoring teams is also important, although hourly monitoring may not be needed, especially if finite-source-size effects increase the time during which the event is observable.

There is another important point as well. This is that the identification of evidence of blending and/or of finite-source-size effects can be important. Because they can possibly clinch the planet-microlensing interpretation, it is worthwhile to search for evidence of these effects in every event. The analysis of light curve shape can play an important role; this analysis can be done after the event has ceased. During the event, it would be valuable to obtain spectra, particularly near the time of peak magnification.

5.1. Useful Modifications

The strategies best suited to the discovery of wide planetary systems have much in common with those best suited to the discovery of planets in resonant orbits. In both cases it is important to have frequent time sampling of events or event features whose duration may be on the order of hours. In both cases, good photometry is important if we are to carry out meaningful tests of the planetary-system lens hypothesis. Thus, since a detailed search mechanism is already in place to discover planets in the zone for resonant lensing, it is useful to ask how effective that mechanism will be at discovering wide planets, and whether slight modifications of it will make it more effective. (See Sackett [1997] for an overview of detection strategies designed for planets in the zone for resonant lensing.)

5.1.1. *Useful Modifications: The Monitoring Teams*

(1) *Spike Analysis:* The MACHO and EROS teams have been able to place upper limits on the numbers of such events along the direction of the LMC (Renault *et al.* 1997, Ansari *et al.* 1996; Alcock *et al.* 1996). Along the direction to the Bulge it seems

likely that the large majority of the events already detected are due to lensing by ordinary stars. If a significant number of stars have planets in wide orbits, then short events due to lensing by these planets are expected. A spike analysis of existing data from the Bulge could be productive in either discovering evidence of short events (in addition to those already discovered) or placing upper limits on the number of short-time-scale events that might have occurred. As more data is collected by a larger number of monitoring teams, systematic spike analyses should become standard.

(2) *Frequent Monitoring of Some Fields:* The importance of the quest for short time-scale events suggests that it might be worthwhile for the teams to each choose one or two fields which they attempt to observe two times per night. This would allow them to call alerts relatively early for Jupiter-mass planets, and to have a better chance of finding evidence of Earth-mass planets. In addition, cooperation among the teams could greatly enhance the probability of finding reliable evidence of short events. At a recent workshop on microlensing, people associated with three of the teams discussed the advantages of choosing one or two fields that all of the teams would attempt to monitor each night. A motivating factor for such cooperation is to better understand the relative detection efficiencies of the teams. The quest for short events provides another important motivation. Indeed, if each team visited one or two Bulge fields twice per night, there would at least occasionally, be 6-8 times per night that those fields were checked. If, on average, 10 events per year were to be discovered in those fields, then, over the course of 3 years, short events could be discovered or ruled out at the $\sim 3\%$ level. In addition, the teams should call alerts for short-duration events, even if the event has apparently ceased before they can announce its discovery. If the short-duration event is due to lensing by a planet, calling the alert will allow the follow-up teams to have a better chance of detecting any subsequent repeat that might be due to lensing by another object in the planetary system.

(3) *Use of Pixel Techniques:* During the past three years, pixel techniques have begun to be used for the study of microlensing in M31 (Tomaney & Crotts 1996; Crotts & Tomaney 1996; Crotts 1996; Ansari *et al.* 1997; Han 1996). It has been estimated that, were such techniques to be applied to the LMC and Bulge fields, the rate of event detection would increase by a factor of $\sim 2 - 3$ (Crotts 1997; Kaplan 1997). The reason for the increase is that observable events can occur when the baseline flux we receive from a star is not bright enough for the star to appear on the templates presently used by the teams, if the flux is brought above the detection limits through microlensing. Such events are presently missed by the monitoring teams. This increase in detection efficiency, would be helpful, particularly in any fields singled out for frequent monitoring. Indeed, the MACHO team is presently engaged in applying pixel subtraction techniques to their LMC data in an attempt to discover events that were missed by their standard methods of detection.

Application of these techniques to the Bulge fields would be also helpful.

5.1.2. *Useful Modifications: The “Follow-Up” Teams*

While monitoring known ongoing events, the follow-up teams have many other stars in their field of view. When the total number, including those not individually above the detection limit (but which could be brought above the limit if magnified by some reasonable amount) is large enough, the follow-up teams can hope to identify new events. Such observations would play a unique role in the identification of new events, particularly the short-duration events that should be associated with wide planetary systems. Programs that would allow the so-called follow-up teams to take the lead in event detection are already underway or are planned (Sahu 1997; Gould 1997). Indeed, an ideal microlensing search for the purposes of the detection of planet lenses, is one in which the follow-up teams play the role now played by the monitoring teams to discover events, and continue the work by monitoring the light curves of the events they discover.

6. Conclusions

Until now, the search for planets via microlensing has focused on a very special detection channel. The work presented here and in the companion paper (DiStefano & Scalzo 1998) strongly argues for an extension of the search to include planets in wide orbits. In this paper we have considered the simplest signal of the presence of planets in wide orbits: isolated events of short-duration, that accompany the passage of the track of a distant star in front of the Einstein ring of an intervening planet. Isolated events of short-duration may be the “vanilla” flavor of planet-lens events. They are nevertheless, interesting for several reasons.

(1) If planetary systems such as our own are common, then, with modest changes in detection strategy, we should discover a few isolated events of short-duration every year. It seems unlikely that our own Solar System is an example of a universal planetary system. It is therefore important to note that a significant fraction of the confirmed planet or brown dwarf systems would also give rise to several, or even tens of isolated short-duration events per year, if *they* were typical of a large fraction of stellar systems.

(2) Isolated events of short-duration may be the most numerous planet-lens events. Even though the detection probability for a planet located in the zone for resonant lensing is enhanced (e.g., for a Jupiter-mass planet, the rate may be as high as 20%), there may

be a large enough number of wide-orbit planets, for every planet in the zone for resonant lensing, to make the detection of isolated events due to planets in wide orbits the dominant mode of planet detection. This is particularly so if (a) sensitive photometry can be employed by the monitoring teams, and (b) finite-source-size effects are important.

(3) Searches can be optimized for the detection of isolated planet-lens events by implementing sensitive photometry. Ideal monitoring programs would use photometry as sensitive as that of the present-day follow-up teams, but would also use difference techniques, in addition to a template, so that they would be sensitive to the lensing of stars too dim to be on the templates. Although frequent monitoring is desirable, hourly monitoring may not be necessary.

(4) Optimized searches can increase the discovery rate significantly. Event rates for systems like our own Solar System, for example, can be doubled. Tests of the hypothesis that all stars are accompanied by a low-mass (Uranus-mass or larger) companion are well within reach and can be carried out over the next few years. Less conservative hypotheses, say that every star is accompanied by a power-of-2 model with Saturn-mass companions, can be ruled out or confirmed in a short time.

(5) If all we know from the data sets is that an ensemble of short-duration events has been observed, then we must rely on statistical arguments (which use the number of stellar-lens events and the numbers of other types of planet-lens events) to determine what fraction of the short-duration events may be due to lensing by planets. Ideally, however, the analysis of individual events could tell us some of the following. (a) the lens is a planet orbiting a luminous star; (b) the spectral type and mass of the star; (c) the mass of the planet; (d) the minimum possible orbital separation between the star and planet. We have pointed out that all of these determinations may be possible for a subset of short-duration events due to planets. This is discussed in more detail elsewhere (DiStefano 1998a,b).

(6) Phenomenologically, overlap events appear to be isolated events, since the associated light curve exhibits just a single deviation from the baseline. Their light curves are distinctive, and provide information about the orbital separation and mass ratio. Overlap events are long-lived relative to isolated events in which just a single planet serves as a lens, and thus they are also relatively easy to discover. The rate of overlap events is determined largely by those planets whose physical location is within a few Einstein radii of the central star. For the Solar System, for example, Jupiter and Saturn are the most likely to contribute to the rate of overlap events. For reasons of dynamical stability, there cannot be a large number of such planets. The ratio of the rate of overlap events relative to the rate of isolated events of short-duration, tells us about the radial structure of planetary systems. In particular it quantifies the size of the population within several R_E of the central star,

relative to the population of planets in wider orbits. This is illustrated by the comparison between the power-of- n models, in which planets populate the system out to $\sim 10^{17}$ cm, and for which the ratio is ~ 0.1 , and the Solar System, where the ratio is close to unity.

Although they may be the most frequent planet-lens events, isolated events of short-duration are only part of the planet-lens story. These events are necessarily accompanied by caustic crossing and repeating events. The statistics and characteristics of planetary systems in other galaxies, and in our own Galactic Bulge, can best be determined by studying the full complement of events. In the accompanying paper, we turn our attention to repeating events.

ACKNOWLEDGEMENTS

We would like to thank the referee, Scott Gaudi, for suggested changes to our original manuscript (DiStefano & Scalzo 1997); his suggestions have helped to improve the presentation. One of us (RD) would like to thank Andrew Becker, Arlin Crotts, Andrew Gould, Jean Kaplan, Christopher Kochanek, David W. Latham, Avi Loeb, Shude Mao, Robert W. Noyes, Bodhan Paczyński, Bill Press, Penny Sackett, Kailash Sahu, Michael M. Shara, Edwin L. Turner, Michael S. Turner, and the participants in the 1997 Aspen workshops, “The Formation and Evolution of Planets” and “Microlensing” for interesting discussions, and the Aspen Center for Physics and the Institute for Theoretical Physics at Santa Barbara for their hospitality while this paper was being written, and the Inter-University Center for Astronomy and Astrophysics in Pune, India for its hospitality while the paper was revised. One of us (RAS) would like to thank the 1996 CfA Summer Intern Program for support and the Harvard-Smithsonian Center for Astrophysics for its hospitality while the work was underway. This work was supported in part by NSF under GER-9450087 and AST-9619516, and by funding from AXAF.

REFERENCES

- Alard, C., Mao, S., & Guibert, J. 1995, A&A, 300L, 17
- Albrow, M., et al. To appear in the Proceedings of the 12th IAP Conference: “Variable Stars and the Astrophysical Returns of Microlensing Surveys”, 1996 (astro-ph/9610128)
- Alcock, C. et al. 1997a, ApJ, 486, 697
- Alcock, C. et al. astro-ph/9702199
- Alcock, C. et al. 1997, ApJ, 479, 119
- Alcock, C. et al. 1996, ApJ, 471, 774
- Alcock, C. et al. 1996, ApJ, 461, 84
- Angel, J. R. P., & Woolf, N. J. 1997, ApJ, 475, 373
- Ansari, R., et al. 1997, A&A, 324, 843
- Ansari, R., et al. 1996, A&A, 314, 94.
- Axelrod 1997, private communication.
- Bender, P. L., & Stebbins, R. T. Journal of Geophysical Research E, 101, 9309
- Bennett, D. P. et al., in “Planets Beyond the Solar System and the Next Generation of Space Missions”, ed. D.R. Soderblom, ASP 119, p 95
- Bennett, D. P. et al. 1996b. Nucl.Phys.Proc.Suppl. 51B (1996) 152-156
- Bennett, D.P. & Rhie, S.H. 1996, ApJ, 472, 660
- Bolatto, A., & Falco, E. 1995, ApJ, 436, 112
- Boutreux, T., & Gould, A. 1996, ApJ, 462, 705
- Bromley, B. C. 1996, ApJ, 467, 537
- Castro, S. *et al.* 1996, AJ, 111, 2439
- Crotts, A. P. S. 1997, private communication
- Crotts, A. P. S. astro-ph/9610067
- Crotts, A. P. S. & Tomaney, A. B. 1996, ApJ, 473, L87

- Di Stefano, R. submitted to ApJ.
- Di Stefano, R. & Scalzo, R.A. 1997, astro-ph/9711013.
- Di Stefano, R. 1998a, submitted to ApJL
- Di Stefano, R. 1998b, in preparation.
- Di Stefano, R., & Mao, S. 1996, ApJ, 457, 93
- Di Stefano, R., & Esin, A. A. 1995, ApJ, 448, L1
- Gaudi, B. S. astro-ph/9706268
- Gaudi, B. S. & Gould, A. 1997, ApJ, 486, 85
- Gould, A. & Welch, D.L., 1996, ApJ, 464, 212
- Gould, A. & Gauchere, C., 1997, ApJ, 477, 580
- Gould, A. 1994, ApJL, 421, 71
- Gould, A. & Loeb, A. 1992, ApJ, 396, 104
- Griest, K. & Hu, W. 1992, ApJ, 397, 362
- Han, C. 1996, ApJ, 472, 108
- Heyrovsky, D. & Loeb, A. 1997, astro-ph/9702097
- Kamionkowski, M. 1995, ApJ, 442, L9
- Kaplan, J. 1997, private communication
- Keeton, C.R. & Kochanek, C.S. 1997, astro-ph/9705194
- Labeyrie, A. 1996, A&AS, 118, 517L
- Latham, D.W. 1997, in “The Origins, Evolutions, and Destinies of Binary Stars in Clusters,”
eds. E.F. Milone and J.-C. Mermilliod, ASP, 90, p 31
- Loeb, A. & Sasselov, D. 1995, ApJ, 449, L33
- Mao, S. & Paczyński, B. 1991, ApJ, 374, L37
- Paczynski et al. 1997, BAAS, 190, 2606

- Paczynski 1996, ARA&A, 34, 419
- Peale, S. J. 1997, Icar, 127, 269
- Peale, S. J. 1997, in “Planets Beyond the Solar System and the Next Generation of Space Missions”, ed. D.R. Soderblom, ASP 119, p 281
- Peng, E.W. 1997, ApJ, 475, 43
- Pratt, M. R., et al. To appear in the Proceedings of the “Source and Detection of Dark Matter in the Universe” Conference held in Santa Monica, CA, Feb., 1996
- Pratt, M. R., et al. Proceedings of the IAU Symposium 173, “Astrophysical Applications of Gravitational Lensing”, Kluwer Academic Publishers, Eds.: C. S. Kochanek and J. N. Hewitt, 1996
- Purdue, P., et al. 1995, AAS, 18710202
- Renault et al. 1997, A&A, 324, 69
- Sackett, P.D. 1997, astro-ph/9709269
- Sahu, K. 1997, in “Planets Beyond the Solar System and the Next Generation of Space Missions”, ed. D.R. Soderblom, ASP 119, p 73
- Sasselov, D. astro-ph/9702180
- Schneider, P., Ehlers, J., & Falco, E.E. 1992, Springer-Verlag
- Simmons, J. F. L., Willis, J. P., & Newsam, A. 1995, A&A, 293, L46
- Stahl, S. M. & Sandler, D. G. 1995, ApJ, 454, L153
- Tomaney, A. B. & Crotts, A. P. S. 1996, AJ, 112, 28
- Udalski, A., Kubiak, M., & Szymanski, M. 1997, Acta Astronomica, 47, 319
- Udalski, A., et al. (astro-ph/9408026)
- Wambsganss, J. 1997, MNRAS, 284, 172
- Witt & Mao, S., 1994, ApJ, 430, 505
- Witt, H.A. 1995, ApJ, 449, 42